



Koppert

TRIANUM[®]-P

Biological Fungicide

ACTIVE INGREDIENT

Trichoderma harzianum Rifai strain T-22* 3.65%

OTHER INGREDIENTS 96.35%

TOTAL 100.00%

*Contains at least 1.0×10^7 colony forming units per g of product.

KEEP OUT OF REACH OF CHILDREN CAUTION

EPA Reg. No. 89635-3

EPA Est. No. 63119-NLD-001

PRODUCT NUMBER 12903

BATCH NUMBER See packaging (10)

NET WEIGHT 1.1 lbs (500 g)

EXPIRATION DATE See Packaging (17) as YYMMDD

MANUFACTURED BY

Koppert Biological Systems, Inc.

1502 Old US 23

Howell, Michigan 48843



400855

v4.0B(5.2)25.01.05



TRIANUM[®]-P

Biological Fungicide

ACTIVE INGREDIENT

<i>Trichoderma harzianum</i> Rifai strain T-22*	3.65%
OTHER INGREDIENTS	96.35%
TOTAL	100.00%

*Contains at least 1.0×10^7 colony forming units per g of product.

KEEP OUT OF REACH OF CHILDREN CAUTION

EPA Reg. No. 89635-3

EPA Est. No. 63119-NLD-001

PRODUCT NUMBER 12903

BATCH NUMBER See packaging (10)

NET WEIGHT 1.1 lbs (500 g)

EXPIRATION DATE See Packaging (17) as YYMMDD

MANUFACTURED BY

Koppert Biological Systems, Inc.
1502 Old US 23
Howell, Michigan 48843



400855

v4.0B(5.2)25.01.05

PRECAUTIONARY STATEMENTS

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Shoes plus socks.

Mixers/loaders and applicators must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R, or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R, or P filter; OR a NIOSH-approved powered air-purifying respirator with an HE filter. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions are available for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected workers may be in the area during application. For any requirement specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water) is protective eyewear, coveralls, waterproof gloves, shoes, and socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants or farms, forests, nurseries or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

PRODUCT INFORMATION

Trianum-P is a preventative control of plant diseases. The active ingredient is a microbe, *Trichoderma harzianum* Rifai strain T-22, which when applied to seeds, transplants or other propagative material, or to soil or planting mixes, grows onto plant roots as they develop and provides protection against plant root pathogens including *Pythium*, *Rhizoctonia*, *Fusarium*, *Cylindrocladium*, *Sclerotinia**, *Sclerotium** and *Thielaviopsis*.

***Not registered for use by California**

Trianum-P is a biological nematicide for control of plant-pest nematodes, including Soybean Cyst Nematode, *Heterodera glycines*.*

***Not registered for use by California**

Trianum-P can be used alone or in conjunction with certain chemical fungicides; consult Trianum-P compatibility chart, tank mix compatibility charts below or the company for more information. This product should not be tank mixed with chemicals that contain the following active ingredients: benzoyl, imazalil, propiconazole, tebuconazole, and triflumizole. Do not apply Trianum-P immediately before these pesticides are used. See specific instructions for tank mixing. Where early season seed rot and seedling diseases are expected, use chemically treated seed or other appropriate measures for stand establishment and Trianum-P for root disease control.

Note: Trianum-P contains live spores of a microbe that must be used prior to disease onset. Trianum-P becomes active in soil or on plants when temperatures are above 50°F and is not effective while temperatures remain cold. Trianum-P can be applied to sterilized or fumigated soil but must be applied after sterilization or fumigation.

This biological fungicide and nematicide* is for use in soil applications (drench, in soil furrow, potting soil and broadcast), and seed treatments in or on all raw agricultural commodities, food and fiber crops. Trianum-P is for use in soil applications (drench, in soil furrow, and potting soil), and seed treatments on ornamentals, landscape plants, turf, and ornamental trees, including tree seedlings for transplanting into the forest.

***Not registered for use by California**

If Trianum-P is mixed with water to make a slurry, dip or suspension, use immediately. If not, keep refrigerated after mixing with water. Refer to the following sections for further mixing instructions:

- Seed Treatment for True Seed Crops.
- Seed Treatment for Vegetatively Propagated Crops, Including Potatoes, other Root, Tuber and Bulb Vegetables.
- Dips for Cuttings and Bare Rooted Transplants.

NOTE: DO NOT APPLY to Sugarcane, Pechay, Rice, Mushrooms, Kiwi, Tobacco, Barley, Oats, Wheat, Lemon, Apple, and Chickpea. Not for use on aquatic crops.

For food commodities: Use in chemigation and irrigation systems is limited to greenhouse flood, drip, furrow, micro-irrigation, and ebb and flow applications with NO OVERHEAD SPRAY. See footnotes for specific directions concerning each use pattern.

APPLY VIA GROUND APPLICATION ONLY

Crops	Use	Rate
Berries and Small Fruits Blackberries, Blueberries, Currants, Elderberries, Gooseberries, Huckleberries, Loganberries, Raspberries, Strawberries, Grapes	Cuttings/bare root	0.25-5.0 lbs / 5 gal or dip into dry powder
	Greenhouse soil drench	1.0-32.0 oz /100 gal
	Nursery soil drench	1.0-32.0 oz /100 gal
	In-furrow spray or transplant starter solution	1.0-32.0 oz / acre
	Greenhouse chemigation ²	1.0-32.0 oz / 100 gal
Bulb Crops Garlic, Leeks, Onions, Shallots, Ornamental Bulbs	Dust (pre-plant)	0.03-3.0 lbs / cwt. seed
Citrus Fruits Citrus Hybrids, Grapefruit, Kumquat, Limes, Oranges, Pummelos	Cuttings or bare-roots	0.25-5.0 lbs / 5 gal or dip into dry powder
	Greenhouse soil drench	1.0-32.0 oz / 100 gal
	Nursery soil drench	1.0-32.0 oz / 100 gal
	In-furrow spray or transplant starter solution	1.0-32.0 oz / acre
	Greenhouse chemigation ²	1.0-32.0 oz / 100 gal
Cucurbit Vegetables Cucumbers, Melons, Gourds, Pumpkins, Squash	Planter Box (on-site)	1.0-10.0 oz / cwt. seed
	Commercial seed treatment ¹	0.00035-26.43 lbs / cwt. seed
	Greenhouse soil drench	1.0-32.0 oz / 100 gal
	In-furrow spray or transplant starter solution	1.0-32.0 oz / acre
	Greenhouse chemigation ²	1.0-32.0 oz / 100 gal
Flowers, Bedding Plants, and Ornamentals	Cuttings or bare-roots	0.25-5.0 lbs / 5 gal or dip into dry powder
	Commercial seed treatment ¹	0.088-17.62 lbs / cwt. seed
	Greenhouse soil drench	1.0-32.0 oz / 100 gal
	Nursery soil drench	1.0-32.0 oz / 100 gal
	Greenhouse chemigation ²	1.0-32.0 oz / 100 gal

Crops	Use	Rate
Fruiting Vegetables Eggplant, Sweet and Hot Peppers, Tomatillos, Tomatoes	Commercial seed treatment ¹ Greenhouse soil drench In-furrow spray or transplant starter solution Greenhouse chemigation ²	0.00035-26.43 lbs / cwt. seed 1.0-32.0 oz / 100 gal 1.0-32.0 oz / acre 1.0-32.0 oz / 100 gal
Herbs, Spices, and Mints	Commercial seed treatment ¹ Greenhouse soil drench In-furrow spray or transplant starter solution Greenhouse chemigation ²	0.00035-26.43 lbs / cwt. seed 1.0-32.0 oz / 100 gal 1.0-32.0 oz / acre 1.0-32.0 oz / 100 gal
Hydroponic Crops Cucumbers, Tomatoes, Lettuce, Herbs and Spices	Greenhouse soil drench Greenhouse chemigation ²	1.0-32.0 oz / 100 gal 1.0-32.0 oz / 100 gal
Leafy Vegetables and Cole Crops Arugula, Celery, Chervil, Endive, Fennel, Lettuce (head and leaf), Parsley, Radicchio, Rhubarb, Spinach, Swiss Chard, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale, Kohlrabi, Mustard Greens, Asparagus	Cuttings or bare-roots Commercial seed treatment ¹ In-furrow spray or transplant starter solution Greenhouse soil drench Greenhouse chemigation ²	0.25-5.0 lbs / 5 gal or dip into dry powder 0.00035 - 26.43 lbs / cwt. seed 1.0-32.0 oz / acre 1.0-32.0 oz / 100 gal 1.0-32.0 oz / 100 gal
Pome Fruit Pears, Quince	Greenhouse soil drench Nursery soil drench In-furrow spray or transplant starter solution Greenhouse chemigation ²	1.0-32.0 oz / 100 gal 1.0-32.0 oz / 100 gal 1.0-32.0 oz / acre 1.0-32.0 oz / 100 gal
Root Crops Beets, Red Beets, Carrots, Celeriac, Chicory, Horseradish, Parsnip, Radish, Rutabaga, Salsify, Turnips	Planter Box (Onsite) Commercial seed treatment ¹ In-furrow spray or transplant starter solution	1.0-10.0 oz / cwt. seed 0.00035-26.43 lbs / cwt. seed 1.0-32.0 oz / acre
Shade house and Outdoor Nursery Crops Deciduous Trees (Maples, Oak, etc.), Ornamentals, Grapes, Citrus, Pine	Cuttings or bare-roots Greenhouse soil drench Nursery soil drench In-furrow spray or transplant starter solution Greenhouse chemigation ²	0.25-5.0 lbs / 5 gal or dip into dry powder 1.0-32.0 oz / 100 gal 1.0-32.0 oz / 100 gal 1.0-32.0 oz / acre 1.0-32.0 oz / 100 gal

Crops	Use	Rate
Stone Fruit Apricots, Cherries, Nectarines, Peaches, Plums, Prunes	Cuttings or bare-roots Greenhouse soil drench Nursery soil drench In-furrow spray or transplant starter solution Greenhouse chemigation ²	0.25-5.0 lbs / 5 gal or dip into dry powder 1.0-32.0 oz / 100 gal 1.0-32.0 oz / 100 gal 1.0-32.0 oz / acre 1.0-32.0 oz / 100 gal
Tree Nuts Almonds, Beech Nuts, Brazil Nuts, Butternuts Cashews, Chestnuts, Filberts, Hickory Nuts, Macadamia Nuts, Pecans, Pistachios, Walnuts	Greenhouse soil drench Nursery soil drench In-furrow spray or transplant starter solution Greenhouse chemigation ²	1.0-32.0 oz / 100 gal 1.0-32.0 oz / 100 gal 1.0-32.0 oz / acre 1.0-32.0 oz / 100 gal

- 1 Refer to Seed Treatment for True Seed Crops section for direction for onsite application to seed.
- 2 Application via greenhouse or field chemigation IS limited to flood, drip, furrow, micro-irrigation, and ebb and flow systems. Do not apply product when above-ground harvestable food commodities are present. Refer to Chemigation section for specific directions.

SEED TREATMENT FOR TRUE SEED CROPS

ONSITE APPLICATION TO SEED

Trium-P is applied to seeds at the rate of 1.0-10.0 oz per hundredweight (oz / cwt.). For example, for large smooth seeds, apply 1.0-10.0 oz / cwt. of seed. For smaller or rougher seed, apply 1.0-8.0 oz / cwt. Trium-P can be applied in sufficient water to coat seeds. For maximum seed protection, especially in cold soils, apply Trium-P to commercially treated seed such as seed treated with Captan, Apron and or Demosan for stand establishment.

Do not use treated seed for food or feed purposes or process for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatments and planting. Do not store excess treated seeds beyond planting time.

COMMERCIAL SEED TREATMENT

Apply Trium-P as slurry, a coating, in a pellet, or during seed priming. See the table on the next page.

NOTE: This product does not contain a dye and is not covered by an appropriate tolerance exemption, or other clearance under the Federal Food, Drug and Cosmetic Act. To comply with 40 CFR §153.155, all seeds treated commercially with this product, must be colored with an EPA-approved colorant of suitable color to prevent accidental use as food for man or feed for animals.

The Federal Seed Act requires that bags containing treated seed shall be labeled with the following information: "This seed has been treated with *Trichoderma harzianum* Rifai strain T-22. Do not use for food, feed or oil purposes."

AGRICULTURAL CROPS

Seed Size (#seeds/oz)	g of Trianum-P per lb of seed	oz of Trianum-P per lb of seed
Large (1-100)	0.0016-0.32 g	0.0016-0.32 oz
Medium (100-1 000)	0.08-2.4 g	0.0028-0.0847 oz
Small (1 000-10 000)	0.4-16 g	0.141-0.564 oz
Fine (10 000-100 000)	1.6-120 g	0.9564-4.23 oz

ORNAMENTAL CROPS

Seed Size (#seeds/oz)	g of Trianum-P per lb of seed	oz of Trianum-P per lb of seed
Small (1 000-10 000) e.g. Echinacea, Cosmos	0.4-16 g	0.0141-0.564 oz
Fine (10 000-100 000) e.g. Texas bluegrass, ryegrass	1.6-120 g	0.0564-4.23 oz
Very fine (100 000-500 000) e.g. bentgrass	8.0-120 g	0.282-2.82 oz

SEED TREATMENT FOR VEGETATIVELY PROPAGATED CROPS, INCLUDING ROOT AND BULB VEGETABLES

For planting or storage, treat at 0.03-3.0 oz Trianum-P to 100 lbs (1 cwt) of bulbs. Dip bulbs in a suspension of 1.0-3.0 lbs of Trianum-P in 20 gal of water.

DIPS FOR CUTTINGS AND BARE ROOTED TRANSPLANTS

Dip cuttings, bulbs or transplants in a suspension of 1.0-3.0 lbs of Trianum-P in 20 gal of water. Plant treated cuttings, bulbs or transplants in potting mix or soil in the usual manner.

SOIL DRENCH

GREENHOUSE SOIL DRENCH

Suspend 1.0-32.0 oz Trianum-P in 100 gal of water with agitation and apply as a soil drench to greenhouse planting mixes. For seeding flats or shallow (up to 4-inch depth) beds or pots, apply at a rate of 50-100 gal per 800 square feet. For deeper beds or pots, apply at a rate of 100 gal per 400 square feet, ½ cup (4 fl. oz) for pots with a 3-inch diameter, or 1 cup (8 fl. oz) per 6-inch diameter pot.

Apply Trianum-P through low pressure watering nozzles such as fan nozzles or other drench watering systems applied directly to the soil. Constant agitation is required to maintain Trianum-P in suspension. Trianum-P can be tank mixed and is compatible with many commonly used fungicides, liquid fertilizers; herbicides, insecticides and biological control products registered for use on greenhouse/ornamental plants.

If tank mixes are desired, observe the most restrictive of labeling limitations and precautions of all products used in mixtures. Consult the tank mix compatibility chart below or the company for more information.

NURSERY SOIL DRENCH

Suspend 1.0-32.0 oz Trianum-P in 100 gal of water with agitation and apply as a soil drench to container nursery crops. For shallow (up to 4-inch depth) beds or pots, apply at a rate of 50-100 gal per 800 square feet. For deeper beds or pots, apply at a rate of 100 gal per 400 square feet, ½ cup (4 fl. oz) for pots with a 3-inch diameter, or 1 cup (8 fl. oz) per 6-inch diameter pot.

Apply Trianum-P directly to the soil through low pressure watering nozzles such as fan nozzles, or other drench watering systems, handheld sprayers or backpack sprayers. Constant agitation is required to maintain Trianum-P in suspension. Trianum-P can be tank mixed and is compatible with many commonly used fungicides, liquid fertilizers, herbicides, insecticides and biological control products registered for use on nursery plants. If tank mixes are desired, observe the most restrictive labeling limitations and precautions of all products used in mixtures. Consult the tank mix compatibility chart below or the company for more information.

IN-FURROW SPRAY OR TRANSPLANT STARTER SOLUTION

Apply as an in-furrow spray or transplant starter solution at a rate of 1.0-32.0 oz/acre in sufficient water to achieve uniform application. Maintain constant agitation. Trianum-P can be tank mixed with certain fertilizers and pesticides; consult tank mix compatibility chart below for detailed information.

TANK MIXING

Trianum-P can be tank mixed and is compatible with many commonly used fungicides, liquid fertilizers, herbicides, insecticides and biological control products registered for use on greenhouse/ornamental plants. If tank mixes are desired, observe the most restrictive of labeling limitations and precautions of all products used in mixtures. Consult the tank mix compatibility chart below or the company for more information.

This product should not be tank mixed with chemicals that contain the following active ingredients: imazilil, propiconazole, tebuconazole, and trilumizole. Do not apply Trianum-P immediately before these pesticides are used. Follow the most restrictive of the labeling limitations and precautions of all products used in mixtures.

This product can be mixed with the specific products, their percentages and rates for use in nursery drench, in-furrow spray or transplant starter solution, as listed in the table below in accordance with the most restrictive of label limitations and precautions. Do not exceed label dosage rates. This product cannot be mixed with any product containing a label prohibition against such mixing.

COMPATIBILITY FOR DRENCH, IN-FURROW SPRAY OR TRANSPLANT STARTER SOLUTION TANK MIX

NOTE: While the information presented in this table is believed to be up to date, the user must always read the label of the other products used in the tank mix to confirm application rates and dilutions.

Chemical Name	% A.I. Formulation	Product Name	Rate Product	Dilution
Captan	85% Wettable Powder	Captan-85WP	1.88 lb / acre	0.1 oz / gal
Chlorothalonil	82.5% Water Dispersible Granules	Daconil Ultrex	3.7 oz / 1 000 ft ²	0.56 oz / gal
Iprodione	23.3% Flowable	Chipco 26019 Flo	4 oz / 1 000 ft ²	0.6 oz / gal
Thiophanate methyl	50% Wettable Powder	Cleary's 3336 in water soluble bags	8 oz / 1 000 ft ²	1.2 oz / gal
Iprodione	Granules 50%, Soluble	Rovral	0.75 lb / acre	0.04 oz / gal
Metalaxyl	21.3% Liquid	SubdueMaxx	0.25 oz / 800 ft ²	0.05 oz / gal
Chlorpyrifos	50% Emulsifiable Liquid	Lorsban4E	3.2 oz / gal	3.2 oz / gal

GREENHOUSE CHEMIGATION

Suspend 1.0-32.0 oz Trium-P in 100 gal of water with agitation and apply through the following systems: 1) pressurized drench (flood) or drip (trickle) systems, 2) furrow, 3) micro-irrigation such as spaghetti-tube or individual tube irrigation, 4) hand-held calibrated irrigation equipment such as the hand-held wand with injector, and 5) ebb and flow systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension specialists, equipment manufacturers or other experts.

Do not connect an irrigation system, (including greenhouse system), used for pesticide application to a public water system unless the pesticide safety systems for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

GREENHOUSE CHEMIGATION

Specific Requirements for Chemigation Systems Connected to Public Water Systems

1. Public water systems means a system for the provision to the public of piped water for human consumption if such a system has at least 15 service connections or regular serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must also contain a functional, normally closed, solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Apply Triatum-P during the last half of the water application period. Mix Triatum-P in enough water to be able to draw through the system for the last half of the water application.
9. Apply enough water to move Triatum-P into the root zone. Amounts will vary depending on soil type and existing moisture level. Avoid applying water volumes that would cause runoff or excessive leaching. Refer to the prior crop tables for specific rate and mixing instructions for application by greenhouse chemigation.

Drip (Trickle) Chemigation and Micro-irrigation Requirements

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Apply Triatum-P during the last half of the water application period. Mix Triatum-P in enough water to be able to draw through the system for the last half of the water application.
8. Apply enough water to move Triatum-P into the root zone. Amounts will vary depending on soil type and existing moisture level. Avoid applying water volumes that would cause runoff or excessive leaching.

PLANT SAFETY

Trianum-P has been tested on numerous plant varieties with no phytotoxic effects. However, since Trianum-P has not been tested on all plant varieties or in combination with all available tank mixes the manufacturer recommends testing Trianum-P on a small number of plants to check for adverse plant effects before applying to a larger number of plants.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE Store in original container under refrigerated conditions. Short periods at room temperatures above 75°F will not affect performance. Do not store near food or feed commodities. Keep container tightly closed when not in use.

PESTICIDE DISPOSAL Wastes resulting from the use of this product must be disposed of on-site or at an approved waste disposal facility.

CONTAINER DISPOSAL Nonrefillable container. Do not reuse or refill this container. Completely empty bag into the application equipment by shaking and tapping sides and bottom to loosen clinging particles, then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration. Do not burn, unless allowed by state and local ordinances. If burned, stay out of smoke.

NOTICE TO BUYER AND SELLER

Seller warrants that this product conforms to the description on the label and is reasonably fit for the purposes stated on the label when used and stored in accordance with directions under normal conditions of use. To the extent permitted by state law, this warranty does not extend to use of this product contrary to label directions or under conditions not reasonably foreseeable by the Seller, and Buyer and User assume the risk of any such use. To the extent permitted by state law, Seller disclaims all other warranties express or implied, including any warranty of fitness or merchantability. To the extent permitted by state law, Seller shall not be liable for consequential, special or indirect damages resulting from use or handling of this product and Seller's sole liability and Buyer's and User's exclusive remedy shall be limited to refund of the purchase price. This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified.

